



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

# ANNALS OF MATHEMATICS.

VOL. IV.

FEBRUARY, 1888.

NO. 1.

## A SOLUTION OF KEPLER'S PROBLEM FOR PLANETARY ORBITS OF HIGH ECCENTRICITY.

By DR. H. A. HOWE, Denver, Col.

The solution of Kepler's Problem involves the transcendental equation

$$M = E - e \sin E, \quad (1)$$

in which  $M$  and  $E$  are respectively the mean and eccentric anomalies, while  $e$  is the eccentricity.

Let

$$\begin{aligned} F(E) &= \frac{1}{2} [E - M - \sin(E - M)] \\ &= \frac{1}{12} \sin^3(E - M) + \frac{3}{80} \sin^5(E - M) \\ &\quad + \frac{5}{224} \sin^7(E - M) + \dots \end{aligned} \quad (2)$$

From (1) and (2) we easily get Grunert's equation,

$$\frac{1 + e}{1 - e} = \frac{\cos \frac{1}{2} M \sin(E - \frac{1}{2} M) + F(E)}{\sin \frac{1}{2} M \cos(E - \frac{1}{2} M) - F(E)}. \quad (3)$$

Equation (3) gives

$$\tan(E - \frac{1}{2} M) = \frac{1 + e}{1 - e} \tan \frac{1}{2} M - \frac{2F(E)}{(1 - e) \cos \frac{1}{2} M \cos(E - \frac{1}{2} M)}. \quad (4)$$

Assume that

$$\tan(E' - \frac{1}{2} M) = \frac{1 + e}{1 - e} \tan \frac{1}{2} M. \quad (5)$$

Substituting in (4) according to (5) and reducing, we obtain

$$\sin(E' - E) = \frac{\cos(E' - \frac{1}{2} M)}{(1 - e) \cos \frac{1}{2} M} \cdot 2F(E) \quad (6)$$

$$= k \cdot 2F(E). \quad (7)$$

Hence for moderate eccentricities,  $E' - E$  is approximately equal to  $2F(E)$ .

Let

$$2\eta = E' - M - \sin(E' - M). \quad (8)$$

Then  $E' - E$  is likewise nearly equal to  $2\eta$ , and  $E' - M - 2\eta$  is nearly equal

to  $E - M$ . From (2) we obtain by differentiation

$$dF(E) = [\frac{1}{4}\sin^2(E - M) + \frac{3}{16}\sin^4(E - M) + \dots] \cos(E - M) d(E - M). \quad (9)$$

Equation (9) shows that a small change in the value of  $E - M$  produces a much smaller change in  $F(E)$ ; therefore, we write with but slight error

$$2F(E) = E' - M - 2\eta - \sin(E' - M - 2\eta). \quad (10)$$

From (8), omitting terms of the seventh and higher orders, we have

$$\eta = \frac{1}{12}\sin^3(E' - M) + \frac{3}{80}\sin^5(E' - M). \quad (11)$$

Suppose that  $E'$  is changed by an amount equal to  $2\eta$ ; i. e. that  $dE' = 2\eta$ . Differentiation of (11) and reduction of the resulting equation, neglecting powers of  $\sin(E' - M)$  higher than the fifth, gives

$$d\eta = \frac{1}{24}\sin^5(E' - M) \cos(E' - M); \quad (12)$$

$$\therefore \eta - d\eta = \frac{1}{12}\sin^3(E' - M) + \frac{3}{80}\sin^5(E' - M) - \frac{1}{24}\sin^5(E' - M) \cos(E' - M). \quad (13)$$

The last two terms of (13) have nearly the same magnitude, and opposite signs.

We therefore write  $\eta - d\eta = \frac{1}{12}\sin^3(E' - M)$ . (14)

Now since (12) is obtained on the assumption that  $dE' = 2\eta$ ,

$$\eta - d\eta = \frac{1}{2}[E' - M - 2\eta - \sin(E' - M - 2\eta)].$$

Hence by (10) and (14) we may write

$$F(E) = \frac{1}{12}\sin^3(E' - M). \quad (15)$$

Equations (15) and (7) give

$$\sin(E' - E) = \frac{1}{6}k \sin^3(E' - M). \quad (16)$$

Let  $E' - M - \frac{1}{6}k \sin^3(E' - M)$  be an approximate value of  $E - M$ ; then  $2F[E' - M - \frac{1}{6}k \sin^3(E' - M)]$  will be the corresponding value of  $2F(E)$ , and by (7),  $k \cdot 2F[E' - M - \frac{1}{6}k \sin^3(E' - M)]$  will be the corresponding value of  $\sin(E' - E)$ . Differentiation of (7) gives

$$\begin{aligned} \cos(E' - E) d(E' - E) &= k [\frac{1}{2}\sin^2(E - M) + \frac{3}{8}\sin^4(E - M) + \dots] \cos(E - M) d(E - M) \\ &= k [\frac{1}{2}\sin^2(E - M) + \frac{1}{8}\sin^4(E - M) + \dots] d(E - M) \\ &= kv d(E - M). \end{aligned} \quad (17)$$

If  $E - M$  be increased by  $E' - E - \frac{1}{6}k \sin^3(E' - M)$ , and the corresponding increment of  $E' - E$  be denoted by  $\Delta(E' - E)$ , we have

$$\Delta(E' - E) = kv [E' - E - \frac{1}{6}k \sin^3(E' - M)] \sec(E' - E). \quad (18)$$

But  $E' - E + \Delta(E' - E) = k \cdot 2F[E' - M - \frac{1}{6}k \sin^3(E' - M)] + 2F(E' - E).$  (19)

Substitute from (18) in (19), and add and subtract  $\frac{1}{6}k \sin^3(E' - M)$  in the second member of the resulting equation; reduction gives

$$(E' - E) [1 + kv \sec(E' - E)] = [1 + kv \sec(E' - E)] \cdot \frac{1}{6}k \sin^3(E' - M) + k \cdot 2F[E' - M - \frac{1}{6}k \sin^3(E' - M)] - \frac{1}{6}k \sin^3(E' - M) + 2F(E' - E). \quad (20)$$

Division of (20) by the coefficient of  $E' - E$ , and subtraction of  $2F(E' - E)$  from each side, gives

$$\begin{aligned} \sin(E' - E) &= \frac{1}{6}k \sin^3(E' - M) \\ &+ \frac{k \{2F[E' - M - \frac{1}{6}k \sin^3(E' - M)] - \frac{1}{6}k \sin^3(E' - M)\}}{1 + kv \sec(E' - E)} \\ &- \frac{kv \sec(E' - E) \cdot 2F(E' - E)}{1 + kv \sec(E' - E)} \end{aligned} \quad (21)$$

But we have, with sufficient accuracy,

$$2F(E' - E) = \frac{1}{6}k \sin^3(E' - E) = \frac{1}{1296}k^3 \sin^9(E' - M),$$

$$\text{and} \quad \frac{kv \sec(E' - E) \cdot 2F(E' - E)}{1 + kv \sec(E' - E)} = \frac{k^4 \sin^{11}(E' - M)}{2592(1 + kv)}. \quad (22)$$

We next determine the error of (21) when its last term is neglected and the quantity  $\sec(E' - E)$  is dropped from the denominator of the second fraction. The eccentricities of the orbits of all but 19 of the first 268 asteroids are below  $\sin 15^\circ$ . Of these 19, 15 lie between  $\sin 15^\circ$  and  $\sin 20^\circ$ , and 4 are greater than  $\sin 20^\circ$ ; the orbit of Aethra (132) has the greatest eccentricity,  $e$  being equal to  $\sin 22^\circ.5$ . On the assumption that  $e = \sin 23^\circ$ , we find that the maximum value of the last term of (21), as computed by (22), is  $0''.003$  +. If the factor  $\sec(E' - E)$  in the denominator of the second fraction of (21) be dropped, an error of  $0''.0002$  may result. To compute  $v$  in the same denominator, we substitute for  $E - M$  (in the expression given for  $v$  in (17)) the value  $E' - M - \frac{1}{6}k \sin^3(E' - M)$ . From equations (17)–(21) we see that, had  $E' - E - \frac{1}{6}k \sin^3(E' - M)$ , the increment of  $E - M$ , been *infinitesimal*,  $E' - M - \frac{1}{6}k \sin^3(E' - M)$  might have been used for  $E - M$  in computing  $v$ , without error. But since  $E' - E - \frac{1}{6}k \sin^3(E' - M)$  is a small finite quantity, the proper value of  $E - M$  to employ is

$$E' - M - \frac{1}{6}k \sin^3(E' - M) - \frac{1}{2}[E' - E - \frac{1}{6}k \sin^3(E' - M)].$$

The greatest error in  $E' - E$  caused by using  $E' - M - \frac{1}{6}k \sin^3(E' - M)$  for  $E - M$ , in finding  $v$ , is less than  $0''.002$ . Greater accuracy is unnecessary, but could be attained, since  $1 + kv$  is always close to unity, by using

$$\begin{aligned} E' - M - \frac{1}{6}k \sin^3(E' - M) \\ - \frac{1}{2}k \{2F[E' - M - \frac{1}{6}k \sin^3(E' - M)] - \frac{1}{6}k \sin^3(E' - M)\}, \end{aligned}$$

or  $E' - M - \frac{1}{12}k \sin^3(E' - M) - kF[E' - M - \frac{1}{6}k \sin^3(E' - M)]$ .

Equation (21) may therefore be written

$$\sin(E' - E) = \frac{1}{6}k \sin^3(E' - M) + \frac{k \{2F[E' - M - \frac{1}{6}k \sin^3(E' - M)] - \frac{1}{6} \sin^3(E' - M)\}}{1 + kv}. \quad (23)$$

When  $e = \sin 23^\circ$ , the maximum numerical value of the last fraction of (23) is  $38''.6$ . A simple approximate value of this fraction in terms of  $k$  and  $\sin(E' - M)$  may be obtained as follows:—

From (9) and (2), by putting  $E' - M$  for  $E - M$ ,  $\frac{1}{6}k \sin^3(E' - M)$  for  $d(E - M)$ , and  $1 - \frac{1}{2}\sin^2(E' - M)$  for  $\cos(E' - M)$ , we get

$$2F[E' - M - \frac{1}{6}k \sin^3(E' - M)] = \frac{1}{6} \sin^3(E' - M) + (\frac{3}{40} - \frac{1}{12}k) \sin^5(E' - M) + (\frac{5}{112} - \frac{1}{48}k) \sin^7(E' - M). \quad (24)$$

Since  $kv$  is small, we assume that

$$\frac{1}{1 + kv} = 1 - kv = 1 - \frac{1}{2}k \sin^2(E' - M). \quad (25)$$

Substitution of (24) and (25) in the last fraction of (23) gives for its value

$$\frac{1}{120}(9k - 10k^2) \sin^5(E' - M) + [\frac{1}{336}(15k - 7k^2) - \frac{1}{240}(9k^2 - 10k^3)] \sin^7(E' - M)$$

Since  $\sin(E' - M) = e \sin E'$ , neglecting powers higher than the fifth, the preceding expression becomes  $\frac{1}{120}(9k - 10k^2) e^5 \sin^5 E'$ . Because this involves the fifth power of  $e$ , its value is small for moderate eccentricities; nevertheless, the expression is not sufficiently accurate to be employed in seven-place computations with large eccentricities. As the second member of (23) is not easy to compute, it may be simplified, and its errors tabulated. It is not difficult to show that the maximum value of  $k$  is  $(1 - e)^{-1}$ , which is reached when  $v = 0$ . The maximum of  $kv$  is near  $\frac{1}{2}e^2$ , which is only  $0.08$  when  $e = \sin 23^\circ$ ;  $kv$  being, therefore, always small, neglect of the denominator of the last term of (23) introduces into  $E' - E$  an error the value of which is not far from

$$\frac{1}{120}kv(9k - 10k^2) e^5 \sin^5 E', \text{ or } \frac{1}{240}(9k^2 - 10k^3) e^7 \sin^7 E'.$$

When  $e = \sin 23^\circ$  this error may amount to  $3''$ ; when  $e = \sin 15^\circ$  it does not reach  $0''.2$ . Rejecting  $1 + kv$  and writing  $E' - E$  for  $\sin(E' - E)$ , (23) becomes

$$E' - E = k \cdot 2F[E' - M - \frac{1}{6}k \sin^3(E' - M)] \quad (26)$$

To expedite computation, two tables are needed, one giving the correction of the second member of (26), the other containing  $2F(x)$  with the argument  $x$ . The computer having  $e$  and  $M$  given, would use (5) and (26), taking into account the tabulated correction to the value of  $E' - E$  given by (26). From the value of  $E$  thus found, the true anomaly could be obtained by one of the usual methods.